




2018 INTEGRATED HIV BEHAVIORAL & SEROLOGIC SURVEILLANCE (IHBSS)

FACTSHEETS



DEPARTMENT OF HEALTH | EPIDEMIOLOGY BUREAU
National HIV/AIDS & STI Surveillance and Strategic
Information Unit (NHSSS)



MALES AND TRANSGENDER WOMEN WHO HAVE SEX WITH MALES - PHILIPPINES

WHO WE SURVEYED

**4098**

MSM & TGW respondents across the Philippines

**53%**

are currently employed

**89%**

lives in the same city of interview


**32%**

are currently studying


Median age (range)

22
(15-67)


AGE GROUP




GENDER IDENTITY



EDUCATIONAL ATTAINMENT



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



RISK BEHAVIORS IN THE PAST 12 MONTHS

**72%**

Had anal sex with a male

**37%**

Received payment in exchange for sex

**31%**

Had sex with a female

**03**

Median number of male sex partners

KNOWLEDGE ON HIV

32%

Know all the five basic facts of HIV transmission and prevention


**82%**
Know that a healthy looking person can have HIV**67%**
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission**80%**
Know that using condoms reduce the risk of HIV transmission**64%**
Know that a person cannot get HIV from mosquito bites**75%**
Know that a person cannot get HIV from using toilet bowls or urinals in public places

Aware that there is a local Social Hygiene Clinic (SHC)



Aware that there is HIV treatment

CONDOM USE AND ACCESS



- NO CONDOM ACCESS
- GETS FREE CONDOMS
- BUYS CONDOMS
- GETS FREE AND BUYS CONDOMS

38%

Condom use during last anal sex (past 12 months)

13%

Consistent condom use (last 3 anal sex partners in the past 12 months)

HIV TESTING AND STI PREVALENCE

43%

Ever tested


32%

Tested in the past 12 months

32%

Know HIV status in the past 12 months*

HIV Hepatitis B Syphilis



*95% confidence interval

IMPORTANT NOTES

Data presented is the aggregate of the 13 cities: Angeles, Baguio, Cagayan De Oro, Cebu, Davao, Gensan, Iloilo, Zamboanga, Mandaue, Talisay, Pasay, Quezon City, Taguig.

-Time location sampling was used.

-Data presented is adjusted using sample weights.

*Tested in the past 12 months and know their status; also includes those who know they are HIV-positive even if their last test was more than 12 months ago

-HIV prevalence is among those who ever had anal sex.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE FEMALE SEX WORKERS PHILIPPINES

WHO WE SURVEYED



2590
FSW respondents



47%
Did sex work all year round



86%
Lives in the same city of interview




92%
Works in an establishment




98%
Works in the same city of interview

Median age (range)
26
(15-56)


AGE GROUP



EDUCATION



CIVIL STATUS



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)

18



First vaginal sex with a male

19



First oral sex with a male

20



First condom use

21



First sex with a male in exchange for cash or kind

22

KNOWLEDGE ON HIV

48%

Know all the five basic facts of HIV transmission and prevention



90%
Know that a healthy looking person can have HIV



79%
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission



88%
Know that using condoms reduce the risk of HIV transmission



73%
Know that a person cannot get HIV from mosquito bites



75%
Know that a person cannot get HIV from using toilet bowls or urinals in public places

45%

Aware that there is HIV treatment

RISK BEHAVIORS



07
Median number of clients in the past 30 days



03
Median number of days with sex work in a week


CONDOM USE AND ACCESS

85%

Condom use with last client

64%

Uses condoms consistently with clients in the past 30 days



NO CONDOM ACCESS
GETS FREE CONDOMS
BUYS CONDOMS
GETS FREE AND BUYS

HIV TESTING AND STI PREVALENCE

72%



Ever tested

57%




Tested in the past 12 months

54%



Tested in the past 12 months and know status

HIV Hepatitis B Syphilis



*95% confidence interval

IMPORTANT NOTES

- Data presented is an aggregate of 10 cities: Angeles, Baguio, Cagayan de Oro, Cebu, Davao, General Santos, Iloilo, Quezon City, Pasay and Zamboanga
- Systematic sampling in health facilities was performed.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph






2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE

MALES AND TRANSGENDER WOMEN WHO HAVE SEX WITH MALES - ANGELES CITY

WHO WE SURVEYED




RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



RISK BEHAVIORS IN THE PAST 12 MONTHS




KNOWLEDGE ON HIV



CONDOM USE AND ACCESS



HIV TESTING AND STI PREVALENCE



IMPORTANT NOTES

- Time location sampling was used.
- Data presented is adjusted using sampling weights.
- * Tested in the past 12 months and know their status; also includes those who know they are HIV-positive even if their last test was more than 12 months ago
- HIV prevalence is among those who ever had anal sex.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE FEMALE SEX WORKERS ANGELES CITY

WHO WE SURVEYED



500
FSW respondents



48%
Did sex work all year round



86%
Lives in the same city of interview




100%
Works in an establishment




100%
Works in the same city of interview

Median age (range)
26
(18-46)


AGE GROUP




EDUCATION



CIVIL STATUS



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



KNOWLEDGE ON HIV

58%

Know all the five basic facts of HIV transmission and prevention



92%
Know that a healthy looking person can have HIV



86%
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission



92%
Know that using condoms reduce the risk of HIV transmission



83%
Know that a person cannot get HIV from mosquito bites



83%
Know that a person cannot get HIV from using toilet bowls or urinals in public places

32%

Aware that there is HIV treatment

RISK BEHAVIORS



04
Median number of clients in the past 30 days



02
Median number of days with sex work in a week


CONDOM USE AND ACCESS

91%

Condom use with last client

72%

Uses condoms consistently with clients in the past 30 days



NO CONDOM ACCESS
GETS FREE CONDOMS
BUYS CONDOMS
GETS FREE AND BUYS

HIV TESTING AND STI PREVALENCE

98%



Ever tested

96%




Tested in the past 12 months

92%



Tested in the past 12 months and know status

HIV Hepatitis B Syphilis



IMPORTANT NOTES

- Systematic sampling in health facilities was performed.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 6517800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





MALES AND TRANSGENDER WOMEN WHO HAVE SEX WITH MALES - BAGUIO CITY

WHO WE SURVEYED



300
MSM & TGW respondents



78%
are currently employed



93%
lives in the same city
of interview




14%
are currently studying


Median age
(range)

25
(15-64)


AGE GROUP




GENDER IDENTITY



EDUCATIONAL ATTAINMENT



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



RISK BEHAVIORS IN THE PAST 12 MONTHS



91%
Had anal sex
with a male



24%
Received payment in
exchange for sex



24%
Had sex with a
female



02
Median number of
male sex partners

KNOWLEDGE ON HIV

52%

Know all the five basic
facts of HIV transmission
and prevention



97%
Know that a healthy
looking person can
have HIV



87%
Know that having sex with
only one, faithful, uninfected
partner can reduce the risk of
HIV transmission



95%
Know that using
condoms reduce the risk
of HIV transmission



67%
Know that a person cannot
get HIV from mosquito
bites



78%
Know that a person cannot get
HIV from using toilet bowls or
urinals in public places




Aware that there is a
local Social Hygiene
Clinic (SHC)



Aware that there is
HIV treatment

CONDOM USE AND ACCESS



● NO CONDOM ACCESS
● GETS FREE CONDOMS
● BUYS CONDOMS
● GETS FREE AND BUYS CONDOMS

33%

Condom use during last anal sex
(past 12 months)

4%

Consistent condom use
(last 3 anal sex partners in the past 12 months)

HIV TESTING AND STI PREVALENCE

38%



Ever tested

20%




Tested in the past
12 months

20%



Know HIV status in
the past 12 months*

HIV Hepatitis B Syphilis



*95% confidence interval

IMPORTANT NOTES

- Time location sampling was used.
- Data presented is adjusted using sampling weights.
* Tested in the past 12 months and know their status; also includes those who know they are HIV-positive even if their last test was more than 12 months ago
- HIV prevalence is among those who ever had anal sex.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE FEMALE SEX WORKERS BAGUIO CITY

WHO WE SURVEYED



200
FSW respondents



41%
Did sex work all year round



99%
Lives in the same city of interview




100%
Works in an establishment




99%
Works in the same city of interview

Median age (range)
29
(18-52)


AGE GROUP



EDUCATION



CIVIL STATUS



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)

18



First vaginal sex with a male

19



First oral sex with a male

20

21

22



First sex with a male in exchange for cash or kind
First condom use

KNOWLEDGE ON HIV

33%

Know all the five basic facts of HIV transmission and prevention



94%
Know that a healthy looking person can have HIV



79%
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission



93%
Know that using condoms reduce the risk of HIV transmission



58%
Know that a person cannot get HIV from mosquito bites



68%
Know that a person cannot get HIV from using toilet bowls or urinals in public places

52%

Aware that there is HIV treatment

RISK BEHAVIORS



03
Median number of clients in the past 30 days



02
Median number of days with sex work in a week


CONDOM USE AND ACCESS

90%

Condom use with last client

72%

Uses condoms consistently with clients in the past 30 days



NO CONDOM ACCESS
GETS FREE CONDOMS
BUYS CONDOMS
GETS FREE AND BUYS

HIV TESTING AND STI PREVALENCE

99%



Ever tested

95%




Tested in the past 12 months

95%



Tested in the past 12 months and know status

HIV Hepatitis B Syphilis



IMPORTANT NOTES

- Systematic sampling in health facilities was performed.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 6517800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE

MALES AND TRANSGENDER WOMEN WHO HAVE SEX WITH MALES - CAGAYAN DE ORO CITY

WHO WE SURVEYED



298

MSM & TGW respondents



60%

are currently employed



90%

lives in the same city of interview




30%

are currently studying


Median age (range)

22
(15-54)


AGE GROUP



GENDER IDENTITY



EDUCATIONAL ATTAINMENT



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



RISK BEHAVIORS IN THE PAST 12 MONTHS



73%

Had anal sex with a male



51%

Received payment in exchange for sex



34%

Had sex with a female



03

Median number of male sex partners

KNOWLEDGE ON HIV

18%

Know all the five basic facts of HIV transmission and prevention



76%

Know that a healthy looking person can have HIV



50%

Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission



58%

Know that using condoms reduce the risk of HIV transmission



55%

Know that a person cannot get HIV from mosquito bites



65%

Know that a person cannot get HIV from using toilet bowls or urinals in public places




Aware that there is a local Social Hygiene Clinic (SHC)



Aware that there is HIV treatment

CONDOM USE AND ACCESS



- NO CONDOM ACCESS
- GETS FREE CONDOMS
- BUYS CONDOMS
- GETS FREE AND BUYS CONDOMS

29%

Condom use during last anal sex (past 12 months)

12%

Consistent condom use (last 3 anal sex partners in the past 12 months)

HIV TESTING AND STI PREVALENCE

29%



Ever tested

19%




Tested in the past 12 months

19%



Know HIV status in the past 12 months*



*95% confidence interval

IMPORTANT NOTES

- Time location sampling was used.
- Data presented is adjusted using sampling weights.
- *Tested in the past 12 months and know their status; also includes those who know they are HIV-positive even if their last test was more than 12 months ago
- HIV prevalence is among those who ever had anal sex.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE FEMALE SEX WORKERS CAGAYAN DE ORO CITY

WHO WE SURVEYED



300
FSW respondents



30%
Did sex work all year round



96%
Lives in the same city of interview




96%
Works in an establishment




97%
Works in the same city of interview

Median age (range)
25
(15-49)


AGE GROUP



EDUCATION



CIVIL STATUS



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)

17



First vaginal sex with a male



18



19



First oral sex with a male



20



First sex with a male in exchange for cash or kind
First condom use



21

KNOWLEDGE ON HIV

31%

Know all the five basic facts of HIV transmission and prevention



89%
Know that a healthy looking person can have HIV



74%
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission



79%
Know that using condoms reduce the risk of HIV transmission



61%
Know that a person cannot get HIV from mosquito bites



65%
Know that a person cannot get HIV from using toilet bowls or urinals in public places

36%

Aware that there is HIV treatment

RISK BEHAVIORS



03
Median number of clients in the past 30 days



02
Median number of days with sex work in a week


CONDOM USE AND ACCESS

71%

Condom use with last client

49%

Uses condoms consistently with clients in the past 30 days



NO CONDOM ACCESS
GETS FREE CONDOMS
BUYS CONDOMS
GETS FREE AND BUYS

HIV TESTING AND STI PREVALENCE

38%



Ever tested

12%




Tested in the past 12 months

11%



Tested in the past 12 months and know status

HIV Hepatitis B Syphilis



*95% confidence interval

IMPORTANT NOTES

- Systematic sampling in health facilities was performed.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 6517800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





MALES AND TRANSGENDER WOMEN WHO HAVE SEX WITH MALES - CEBU CITY

WHO WE SURVEYED

**459**

MSM & TGW respondents

**40%**

are currently employed

**90%**

lives in the same city of interview


**32%**

are currently studying


Median age (range)

22
(15-57)


AGE GROUP




GENDER IDENTITY



EDUCATIONAL ATTAINMENT



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



RISK BEHAVIORS IN THE PAST 12 MONTHS

**77%**

Had anal sex with a male

**44%**

Received payment in exchange for sex

**21%**

Had sex with a female

**05**

Median number of male sex partners

KNOWLEDGE ON HIV

26%

Know all the five basic facts of HIV transmission and prevention

**83%**

Know that a healthy looking person can have HIV

**63%**

Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission

**78%**

Know that using condoms reduce the risk of HIV transmission

**66%**

Know that a person cannot get HIV from mosquito bites

**69%**

Know that a person cannot get HIV from using toilet bowls or urinals in public places




Aware that there is a local Social Hygiene Clinic (SHC)



Aware that there is HIV treatment

CONDOM USE AND ACCESS



- NO CONDOM ACCESS
- GETS FREE CONDOMS
- BUYS CONDOMS
- GETS FREE AND BUYS CONDOMS

42%

Condom use during last anal sex (past 12 months)

13%

Consistent condom use (last 3 anal sex partners in the past 12 months)

HIV TESTING AND STI PREVALENCE

54%

Ever tested


40%

Tested in the past 12 months

39%

Know HIV status in the past 12 months*

HIV Hepatitis B Syphilis Hepatitis C



*95% confidence interval

IMPORTANT NOTES

- Time location sampling was used.
- Data presented is adjusted using sampling weights.
- * Tested in the past 12 months and know their status; also includes those who know they are HIV-positive even if their last test was more than 12 months ago
- HIV prevalence is among those who ever had anal sex.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE FEMALE SEX WORKERS CEBU CITY

WHO WE SURVEYED



200
FSW respondents



59%
Did sex work all year round



73%
Lives in the same city of interview




98%
Works in an establishment




96%
Works in the same city of interview

Median age (range)
28
(18-51)


AGE GROUP



EDUCATION



CIVIL STATUS



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)

18



First vaginal sex with a male

19

20



First oral sex with a male

21



First sex with a male in exchange for cash or kind
First condom use

22

KNOWLEDGE ON HIV

55%

Know all the five basic facts of HIV transmission and prevention



84%
Know that a healthy looking person can have HIV



86%
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission



88%
Know that using condoms reduce the risk of HIV transmission



77%
Know that a person cannot get HIV from mosquito bites



78%
Know that a person cannot get HIV from using toilet bowls or urinals in public places

37%

Aware that there is HIV treatment

RISK BEHAVIORS



05
Median number of clients in the past 30 days



02
Median number of days with sex work in a week


CONDOM USE AND ACCESS

92%

Condom use with last client

79%

Uses condoms consistently with clients in the past 30 days



NO CONDOM ACCESS
GETS FREE CONDOMS
BUYS CONDOMS
GETS FREE AND BUYS

HIV TESTING AND STI PREVALENCE

95%



Ever tested

85%




Tested in the past 12 months

83%



Tested in the past 12 months and know status

HIV Hepatitis B Hepatitis C Syphilis



IMPORTANT NOTES

- Systematic sampling in health facilities was performed.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 6517800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





MALES AND TRANSGENDER WOMEN WHO HAVE SEX WITH MALES - DAVAO CITY

WHO WE SURVEYED



301
MSM & TGW respondents



63%
are currently employed



99%
lives in the same city
of interview




25%
are currently studying


Median age
(range)

22
(15-54)


AGE GROUP




GENDER IDENTITY



EDUCATIONAL ATTAINMENT



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



RISK BEHAVIORS IN THE PAST 12 MONTHS



71%
Had anal sex
with a male



28%
Received payment in
exchange for sex



16%
Had sex with a
female



03
Median number of
male sex partners

KNOWLEDGE ON HIV

41%

Know all the five basic
facts of HIV transmission
and prevention



92%
Know that a healthy
looking person can
have HIV



73%
Know that having sex with
only one, faithful, uninfected
partner can reduce the risk of
HIV transmission



80%
Know that using
condoms reduce the risk of
HIV transmission



72%
Know that a person cannot
get HIV from mosquito
bites



89%
Know that a person cannot get
HIV from using toilet bowls or
urinals in public places




Aware that there is a
local Social Hygiene
Clinic (SHC)



Aware that there is
HIV treatment

CONDOM USE AND ACCESS



- NO CONDOM ACCESS
- GETS FREE CONDOMS
- BUYS CONDOMS
- GETS FREE AND BUYS CONDOMS

49%

Condom use during last anal sex
(past 12 months)

14%

Consistent condom use
(last 3 anal sex partners in the past 12 months)

HIV TESTING AND STI PREVALENCE

62%



Ever tested

42%




Tested in the past
12 months

45%



Know HIV status in
the past 12 months *



IMPORTANT NOTES

- Time location sampling was used.
- Data presented is adjusted using sampling weights.
- * Tested in the past 12 months and know their status; also includes those who know they are HIV-positive even if their last test was more than 12 months ago
- HIV prevalence is among those who ever had anal sex.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE FEMALE SEX WORKERS DAVAO CITY

WHO WE SURVEYED



212
FSW respondents



56%
Did sex work all year round



98%
Lives in the same city of interview




78%
Works in an establishment




97%
Works in the same city of interview

Median age (range)
27
(18-56)


AGE GROUP



EDUCATION



CIVIL STATUS



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)

17



First vaginal sex with a male

18

19

20



First oral sex with a male
First condom use

21



First sex with a male in exchange for cash or kind

KNOWLEDGE ON HIV

60%

Know all the five basic facts of HIV transmission and prevention



87%
Know that a healthy looking person can have HIV



85%
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission



91%
Know that using condoms reduce the risk of HIV transmission



73%
Know that a person cannot get HIV from mosquito bites



81%
Know that a person cannot get HIV from using toilet bowls or urinals in public places



Aware that there is HIV treatment

RISK BEHAVIORS



04
Median number of clients in the past 30 days



02
Median number of days with sex work in a week


CONDOM USE AND ACCESS



Condom use with last client



Uses condoms consistently with clients in the past 30 days



NO CONDOM ACCESS
GETS FREE CONDOMS
BUYS CONDOMS
GETS FREE AND BUYS

HIV TESTING AND STI PREVALENCE

76%



Ever tested

50%




Tested in the past 12 months

42%



Tested in the past 12 months and know status

HIV Hepatitis B Syphilis



IMPORTANT NOTES

- Systematic sampling in health facilities was performed.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 6517800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE

MALES AND TRANSGENDER WOMEN WHO HAVE SEX WITH MALES - GENERAL SANTOS CITY

WHO WE SURVEYED



299

MSM & TGW respondents



52%

are currently employed



97%

lives in the same city of interview




36%

are currently studying


Median age (range)

22
(15-52)


AGE GROUP




GENDER IDENTITY



EDUCATIONAL ATTAINMENT



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



RISK BEHAVIORS IN THE PAST 12 MONTHS



57%

Had anal sex with a male



47%

Received payment in exchange for sex



35%

Had sex with a female



02

Median number of male sex partners

KNOWLEDGE ON HIV

19%

Know all the five basic facts of HIV transmission and prevention



65%
Know that a healthy looking person can have HIV



63%
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission



74%
Know that using condoms reduce the risk of HIV transmission



53%
Know that a person cannot get HIV from mosquito bites



72%
Know that a person cannot get HIV from using toilet bowls or urinals in public places




Aware that there is a local Social Hygiene Clinic (SHC)



Aware that there is HIV treatment

CONDOM USE AND ACCESS



- NO CONDOM ACCESS
- GETS FREE CONDOMS
- BUYS CONDOMS
- GETS FREE AND BUYS CONDOMS

34%

Condom use during last anal sex (past 12 months)

2%

Consistent condom use (last 3 anal sex partners in the past 12 months)

HIV TESTING AND STI PREVALENCE

16%



Ever tested

4%




Tested in the past 12 months

3%



Know HIV status in the past 12 months*

HIV Hepatitis B Syphilis



IMPORTANT NOTES

- Time location sampling was used.
- Data presented is adjusted using sampling weights.
- * Tested in the past 12 months and know their status; also includes those who know they are HIV-positive even if their last test was more than 12 months ago
- HIV prevalence is among those who ever had anal sex.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE FEMALE SEX WORKERS GENERAL SANTOS CITY

WHO WE SURVEYED



150
FSW respondents



67%
Did sex work all year round



97%
Lives in the same city of interview




87%
Works in an establishment




97%
Works in the same city of interview

Median age (range)
27
(15-54)


AGE GROUP



EDUCATION



CIVIL STATUS



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)

17



First vaginal sex with a male

18

19

20



First oral sex with a male
First condom use
First sex with a male in exchange for cash or kind

21

KNOWLEDGE ON HIV

54%

Know all the five basic facts of HIV transmission and prevention



93%
Know that a healthy looking person can have HIV



75%
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission



87%
Know that using condoms reduce the risk of HIV transmission



83%
Know that a person cannot get HIV from mosquito bites



66%
Know that a person cannot get HIV from using toilet bowls or urinals in public places

81%

Aware that there is HIV treatment

RISK BEHAVIORS



10
Median number of clients in the past 30 days



03
Median number of days with sex work in a week


CONDOM USE AND ACCESS

91%

Condom use with last client

85%

Uses condoms consistently with clients in the past 30 days



NO CONDOM ACCESS
GETS FREE CONDOMS
BUYS CONDOMS
GETS FREE AND BUYS

HIV TESTING AND STI PREVALENCE

56%



Ever tested

42%




Tested in the past 12 months

37%



Tested in the past 12 months and know status

HIV Hepatitis B Syphilis



IMPORTANT NOTES

- Systematic sampling in health facilities was performed.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 6517800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





MALES AND TRANSGENDER WOMEN WHO HAVE SEX WITH MALES - ILOILO CITY

WHO WE SURVEYED

**303**

MSM & TGW respondents

**54%**

are currently employed

**98%**

lives in the same city of interview


**34%**

are currently studying


Median age (range)

24
(15-67)


AGE GROUP



GENDER IDENTITY



EDUCATIONAL ATTAINMENT



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)

15First oral sex with a male
First anal sex with a male**17**

First sex with a female

16**18**

First condom use

19**20**

First HIV test

21

RISK BEHAVIORS IN THE PAST 12 MONTHS

**71%**

Had anal sex with a male

**37%**

Received payment in exchange for sex

**20%**

Had sex with a female

**03**

Median number of male sex partners

KNOWLEDGE ON HIV

23%

Know all the five basic facts of HIV transmission and prevention


**72%**
Know that a healthy looking person can have HIV**68%**
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission**79%**
Know that using condoms reduce the risk of HIV transmission**52%**
Know that a person cannot get HIV from mosquito bites**71%**
Know that a person cannot get HIV from using toilet bowls or urinals in public places

Aware that there is a local Social Hygiene Clinic (SHC)



Aware that there is HIV treatment

CONDOM USE AND ACCESS



- NO CONDOM ACCESS
- GETS FREE CONDOMS
- BUYS CONDOMS
- GETS FREE AND BUYS CONDOMS

50%

Condom use during last anal sex (past 12 months)

30%

Consistent condom use (last 3 anal sex partners in the past 12 months)

HIV TESTING AND STI PREVALENCE

52%

Ever tested

37%

Tested in the past 12 months


36%

Know HIV status in the past 12 months*

HIV

Hepatitis B

Syphilis



*95% confidence interval

IMPORTANT NOTES

- Time location sampling was used.
- Data presented is adjusted using sampling weights.
- * Tested in the past 12 months and know their status; also includes those who know they are HIV-positive even if their last test was more than 12 months ago
- HIV prevalence is among those who ever had anal sex.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE FEMALE SEX WORKERS ILOILO CITY

WHO WE SURVEYED



306
FSW respondents



40%
Did sex work all year round



96%
Lives in the same city of interview




64%
Works in an establishment




96%
Works in the same city of interview

Median age (range)
27
(18-55)


AGE GROUP



EDUCATION



CIVIL STATUS



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)

17



First vaginal sex with a male

18

19

20



First oral sex with a male

21



First condom use
First sex with a male in exchange for cash or kind

KNOWLEDGE ON HIV

42%

Know all the five basic facts of HIV transmission and prevention



89%
Know that a healthy looking person can have HIV



71%
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission



86%
Know that using condoms reduce the risk of HIV transmission



72%
Know that a person cannot get HIV from mosquito bites



84%
Know that a person cannot get HIV from using toilet bowls or urinals in public places

24%

Aware that there is HIV treatment

RISK BEHAVIORS



15
Median number of clients in the past 30 days



03
Median number of days with sex work in a week


CONDOM USE AND ACCESS

83%

Condom use with last client

66%

Uses condoms consistently with clients in the past 30 days



NO CONDOM ACCESS
GETS FREE CONDOMS
BUYS CONDOMS
GETS FREE AND BUYS

HIV TESTING AND STI PREVALENCE

45%



Ever tested

23%




Tested in the past 12 months

22%



Tested in the past 12 months and know status

HIV Hepatitis B Syphilis



*95% confidence interval

IMPORTANT NOTES

- Systematic sampling in health facilities was performed.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 6517800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE

MALES AND TRANSGENDER WOMEN WHO HAVE SEX WITH MALES - MANDAUE CITY

WHO WE SURVEYED



306
MSM & TGW respondents



56%
are currently employed



87%
lives in the same city
of interview




30%
are currently studying


Median age
(range)

22
(15-56)


AGE GROUP




GENDER IDENTITY



EDUCATIONAL ATTAINMENT



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



RISK BEHAVIORS IN THE PAST 12 MONTHS



93%
Had anal sex
with a male



47%
Received payment in
exchange for sex



65%
Had sex with a
female



06
Median number of
male sex partners

KNOWLEDGE ON HIV

15%

Know all the five basic
facts of HIV transmission
and prevention



76%
Know that a healthy
looking person can
have HIV



50%
Know that having sex with
only one, faithful, uninfected
partner can reduce the risk of
HIV transmission



78%
Know that using
condoms reduce the risk of
HIV transmission



56%
Know that a person cannot
get HIV from mosquito
bites



67%
Know that a person cannot get
HIV from using toilet bowls or
urinals in public places




Aware that there is a
local Social Hygiene
Clinic (SHC)



Aware that there is
HIV treatment

CONDOM USE AND ACCESS



- NO CONDOM ACCESS
- GETS FREE CONDOMS
- BUYS CONDOMS
- GETS FREE AND BUYS CONDOMS

42%

Condom use during last anal sex
(past 12 months)

13%

Consistent condom use
(last 3 anal sex partners in the past 12 months)

HIV TESTING AND STI PREVALENCE

28%



Ever tested

24%




Tested in the past
12 months

24%



Know HIV status in
the past 12 months*

HIV Hepatitis B Syphilis Hepatitis C



*95% confidence interval

IMPORTANT NOTES

- Time location sampling was used.
- Data presented is adjusted using sampling weights.
- * Tested in the past 12 months and know their status; also includes those who know they are HIV-positive even if their last test was more than 12 months ago
- HIV prevalence is among those who ever had anal sex.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





MALES AND TRANSGENDER WOMEN WHO HAVE SEX WITH MALES - PASAY CITY

WHO WE SURVEYED

**301**

MSM & TGW respondents

**44%**

are currently employed

**77%**

lives in the same city of interview


**53%**

are currently studying


Median age (range)

21
(15-58)


AGE GROUP



GENDER IDENTITY



EDUCATIONAL ATTAINMENT



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)

15**17**First anal sex with a male
First condom use**19****21****16**First sex with a female
First oral sex with a male**18****20**

First HIV test

RISK BEHAVIORS IN THE PAST 12 MONTHS

**53%**

Had anal sex with a male

**34%**

Received payment in exchange for sex

**46%**

Had sex with a female

**02**

Median number of male sex partners

KNOWLEDGE ON HIV

33%

Know all the five basic facts of HIV transmission and prevention


**87%**
Know that a healthy looking person can have HIV**72%**
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission**82%**
Know that using condoms reduce the risk of HIV transmission**66%**
Know that a person cannot get HIV from mosquito bites**74%**
Know that a person cannot get HIV from using toilet bowls or urinals in public places

Aware that there is a local Social Hygiene Clinic (SHC)



Aware that there is HIV treatment

CONDOM USE AND ACCESS



- NO CONDOM ACCESS
- GETS FREE CONDOMS
- BUYS CONDOMS
- GETS FREE AND BUYS CONDOMS

43%

Condom use during last anal sex (past 12 months)

10%

Consistent condom use (last 3 anal sex partners in the past 12 months)

HIV TESTING AND STI PREVALENCE

33%

Ever tested


23%

Tested in the past 12 months

21%

Know HIV status in the past 12 months*

HIV Hepatitis B Syphilis



*95% confidence interval

IMPORTANT NOTES

- Time location sampling was used.
- Data presented is adjusted using sampling weights.
- * Tested in the past 12 months and know their status; also includes those who know they are HIV-positive even if their last test was more than 12 months ago
- HIV prevalence is among those who ever had anal sex.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE FEMALE SEX WORKERS PASAY CITY

WHO WE SURVEYED



150

FSW respondents



33%

Did sex work all year round



36%

Lives in the same city of interview



100%

Works in an establishment




99%


Works in the same city of interview

Median age (range)
25
(18-45)


AGE GROUP



EDUCATION



CIVIL STATUS



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)

17



First vaginal sex with a male

18

19



First oral sex with a male

20



First condom use

21



First sex with a male in exchange for cash or kind

KNOWLEDGE ON HIV

17%

Know all the five basic facts of HIV transmission and prevention



86%

Know that a healthy looking person can have HIV



67%

Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission



86%

Know that using condoms reduce the risk of HIV transmission




50%

Know that a person cannot get HIV from mosquito bites



47%

Know that a person cannot get HIV from using toilet bowls or urinals in public places



Aware that there is HIV treatment

RISK BEHAVIORS



10

Median number of clients in the past 30 days



03

Median number of days with sex work in a week


CONDOM USE AND ACCESS



Condom use with last client



Uses condoms consistently with clients in the past 30 days



NO CONDOM ACCESS
GETS FREE CONDOMS
BUYS CONDOMS
GETS FREE AND BUYS

HIV TESTING AND STI PREVALENCE

33%



Ever tested

6%



Tested in the past 12 months

5%



Tested in the past 12 months and know status

HIV Hepatitis B Syphilis



IMPORTANT NOTES

- Systematic sampling in health facilities was performed.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 6517800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





MALES AND TRANSGENDER WOMEN WHO HAVE SEX WITH MALES - QUEZON CITY

WHO WE SURVEYED

**427**

MSM & TGW respondents

**61%**

are currently employed

**69%**

lives in the same city of interview


**23%**

are currently studying


Median age (range)

24
(15-67)


AGE GROUP




GENDER IDENTITY



EDUCATIONAL ATTAINMENT



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



RISK BEHAVIORS IN THE PAST 12 MONTHS

**60%**

Had anal sex with a male

**27%**

Received payment in exchange for sex

**28%**

Had sex with a female

**03**

Median number of male sex partners

KNOWLEDGE ON HIV

50%

Know all the five basic facts of HIV transmission and prevention

**92%**

Know that a healthy looking person can have HIV

**81%**

Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV

**91%**

Know that using condoms reduce the risk of HIV transmission

**69%**

Know that a person cannot get HIV from mosquito bites

**85%**

Know that a person cannot get HIV from using toilet bowls or urinals in public places




Aware that there is a local Social Hygiene Clinic (SHC)



Aware that there is HIV treatment

CONDOM USE AND ACCESS



- NO CONDOM ACCESS
- GETS FREE CONDOMS
- BUYS CONDOMS
- GETS FREE AND BUYS CONDOMS

38%

Condom use during last anal sex (past 12 months)

15%

Consistent condom use (last 3 anal sex partners in the past 12 months)

HIV TESTING AND STI PREVALENCE

53%


Ever tested

44%

Tested in the past 12 months

42%

Know HIV status in the past 12 months*



IMPORTANT NOTES

- Time location sampling was used.
- Data presented is adjusted using sampling weights.
- * Tested in the past 12 months and know their status; also includes those who know they are HIV-positive even if their last test was more than 12 months ago
- HIV prevalence is among those who ever had anal sex.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE FEMALE SEX WORKERS QUEZON CITY

WHO WE SURVEYED



322
FSW respondents



35%
Did sex work all year round



65%
Lives in the same city of interview




98%
Works in an establishment




97%
Works in the same city of interview

Median age (range)
25
(18-45)


AGE GROUP




EDUCATION



CIVIL STATUS



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



KNOWLEDGE ON HIV

44%

Know all the five basic facts of HIV transmission and prevention



87%
Know that a healthy looking person can have HIV



74%
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission



84%
Know that using condoms reduce the risk of HIV transmission



77%
Know that a person cannot get HIV from mosquito bites



76%
Know that a person cannot get HIV from using toilet bowls or urinals in public places



Aware that there is HIV treatment

RISK BEHAVIORS



10
Median number of clients in the past 30 days



03
Median number of days with sex work in a week


CONDOM USE AND ACCESS



Condom use with last client



Uses condoms consistently with clients in the past 30 days



NO CONDOM ACCESS
GETS FREE CONDOMS
BUYS CONDOMS
GETS FREE AND BUYS

HIV TESTING AND STI PREVALENCE

80%



Ever tested

69%




Tested in the past 12 months

66%



Tested in the past 12 months and know status

HIV Hepatitis B Syphilis



IMPORTANT NOTES

- Systematic sampling in health facilities was performed.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 6517800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





MALES AND TRANSGENDER WOMEN WHO HAVE SEX WITH MALES - TAGUIG CITY

WHO WE SURVEYED



310
MSM & TGW respondents



50%
are currently employed



85%
lives in the same city
of interview




24%
are currently studying


Median age
(range)

22
(15-51)


AGE GROUP




GENDER IDENTITY



EDUCATIONAL ATTAINMENT



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



RISK BEHAVIORS IN THE PAST 12 MONTHS



71%
Had anal sex
with a male



6%
Had sex with a male in
exchange for cash or
kind



18%
Had sex with a
female



03
Median number of
male sex partners

KNOWLEDGE ON HIV

57%

Know all the five basic
facts of HIV transmission
and prevention



93%
Know that a healthy
looking person can
have HIV



77%
Know that having sex with
only one, faithful, uninfected
partner can reduce the risk of
HIV transmission



93%
Know that using
condoms reduce the risk of
HIV transmission



83%
Know that a person cannot
get HIV from mosquito
bites



87%
Know that a person cannot get
HIV from using toilet bowls or
urinals in public places




Aware that there is a
local Social Hygiene
Clinic (SHC)



Aware that there is
HIV treatment

CONDOM USE AND ACCESS



- NO CONDOM ACCESS
- GETS FREE CONDOMS
- BUYS CONDOMS
- GETS FREE AND BUYS CONDOMS

40%

Condom use during last anal sex
(past 12 months)

13%

Consistent condom use
(last 3 anal sex partners in the past 12 months)

HIV TESTING AND STI PREVALENCE

68%



Ever tested

61%




Tested in the past
12 months

60%



Know HIV status in
the past 12 months*

HIV Hepatitis B Syphilis



*95% confidence interval

IMPORTANT NOTES

- Time location sampling was used.
- Data presented is adjusted using sampling weights.
- * Tested in the past 12 months and know their status; also includes those who know they are HIV-positive even if their last test was more than 12 months ago
- HIV prevalence is among those who ever had anal sex.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





MALES AND TRANSGENDER WOMEN WHO HAVE SEX WITH MALES - TALISAY CITY, CEBU

WHO WE SURVEYED



300
MSM & TGW respondents



29%
are currently employed



97%
lives in the same city
of interview




54%
are currently studying


Median age
(range)

19
(15-59)


AGE GROUP




GENDER IDENTITY



EDUCATIONAL ATTAINMENT



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



RISK BEHAVIORS IN THE PAST 12 MONTHS



93%
Had anal sex
with a male



56%
Received payment in
exchange for sex



32%
Had sex with a
female



03
Median number of
male sex partners

KNOWLEDGE ON HIV

28%

Know all the five basic
facts of HIV transmission
and prevention



73%
Know that a healthy
looking person can
have HIV



57%
Know that having sex with
only one, faithful, uninfected
partner can reduce the risk of
HIV transmission



74%
Know that using
condoms reduce the risk of
HIV transmission



63%
Know that a person cannot
get HIV from mosquito
bites



75%
Know that a person cannot get
HIV from using toilet bowls or
urinals in public places




Aware that there is a
local Social Hygiene
Clinic (SHC)



Aware that there is
HIV treatment

CONDOM USE AND ACCESS



- NO CONDOM ACCESS
- GETS FREE CONDOMS
- BUYS CONDOMS
- GETS FREE AND BUYS CONDOMS

21%

Condom use during last anal sex
(past 12 months)

7%

Consistent condom use
(last 3 anal sex partners in the past 12 months)

HIV TESTING AND STI PREVALENCE

43%



Ever tested

39%



Tested in the past
12 months

37%



Know HIV status in
the past 12 months*

HIV Hepatitis B Syphilis



*95% confidence interval

IMPORTANT NOTES

- Time location sampling was used.
- Data presented is adjusted using sampling weights.
- * Tested in the past 12 months and know their status; also includes those who know they are HIV-positive even if their last test was more than 12 months ago
- HIV prevalence is among those who ever had anal sex.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





MALES AND TRANSGENDER WOMEN WHO HAVE SEX WITH MALES - ZAMBOANGA CITY

WHO WE SURVEYED

**300**

MSM & TGW respondents

**53%**

are currently employed

**100%**

lives in the same city of interview


**28%**

are currently studying


Median age (range)

22
(15-65)


AGE GROUP



GENDER IDENTITY



EDUCATIONAL ATTAINMENT



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)



RISK BEHAVIORS IN THE PAST 12 MONTHS

**60%**

Had anal sex with a male

**40%**

Received payment in exchange for sex

**35%**

Had sex with a female

**01**

Median number of male sex partners

KNOWLEDGE ON HIV

27%

Know all the five basic facts of HIV transmission and prevention


**73%**
Know that a healthy looking person can have HIV**57%**
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission**75%**
Know that using condoms reduce the risk of HIV transmission**63%**
Know that a person cannot get HIV from mosquito bites**69%**
Know that a person cannot get HIV from using toilet bowls or urinals in public places**95%**

Aware that there is a local Social Hygiene Clinic (SHC)

10%

Aware that there is HIV treatment

CONDOM USE AND ACCESS



- NO CONDOM ACCESS
- GETS FREE CONDOMS
- BUYS CONDOMS
- GETS FREE AND BUYS CONDOMS

20%

Condom use during last anal sex (past 12 months)

6%

Consistent condom use (last 3 anal sex partners in the past 12 months)

HIV TESTING AND STI PREVALENCE

25%

Ever tested


19%

Tested in the past 12 months

19%

Know HIV status in the past 12 months*

HIV Hepatitis B Syphilis



*95% confidence interval

IMPORTANT NOTES

- Time location sampling was used.
- Data presented is adjusted using sampling weights.
- * Tested in the past 12 months and know their status; also includes those who know they are HIV-positive even if their last test was more than 12 months ago
- HIV prevalence is among those who ever had anal sex.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 651-7800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph





2018 PHILIPPINE INTEGRATED HIV BEHAVIORAL AND SEROLOGIC SURVEILLANCE FEMALE SEX WORKERS ZAMBOANGA CITY

WHO WE SURVEYED



250
FSW respondents



74%
Did sex work all year round



100%
Lives in the same city of interview




97%
Works in an establishment




100%
Works in the same city of interview

Median age (range)
26
(18-53)


AGE GROUP



EDUCATION



CIVIL STATUS



RISK AND PROTECTIVE BEHAVIOR MILESTONES (MEDIAN AGE)

17



First vaginal sex with a male

18



First oral sex with a male

19



First sex with a male in exchange for cash or kind

20



First condom use

21

KNOWLEDGE ON HIV

70%

Know all the five basic facts of HIV transmission and prevention



92%
Know that a healthy looking person can have HIV



86%
Know that having sex with only one, faithful, uninfected partner can reduce the risk of HIV transmission



93%
Know that using condoms reduce the risk of HIV transmission



82%
Know that a person cannot get HIV from mosquito bites



84%
Know that a person cannot get HIV from using toilet bowls or urinals in public places

62%

Aware that there is HIV treatment

RISK BEHAVIORS



20
Median number of clients in the past 30 days



05
Median number of days with sex work in a week


CONDOM USE AND ACCESS

76%

Condom use with last client

25%

Uses condoms consistently with clients in the past 30 days



HIV TESTING AND STI PREVALENCE

74%



Ever tested

52%




Tested in the past 12 months

52%



Tested in the past 12 months and know status

HIV Hepatitis B Syphilis



IMPORTANT NOTES

- Systematic sampling in health facilities was performed.



NATIONAL HIV/AIDS & STI SURVEILLANCE AND STRATEGIC INFORMATION UNIT

Epidemiology Bureau, Department of Health
Bldg. 19, 2nd Floor, San Lazaro Compound,
Sta. Cruz, Manila, 1003, Philippines

Tel: +632 6517800 loc 2952
Email: hivpicenter@gmail.com
Website: www.doh.gov.ph

