



Measles Elimination Goal in the Philippines

Measles elimination goal is the absence of endemic measles virus transmission in a defined geographical area (e.g. region or country) for at least 12 months in the presence of a surveillance system that has been verified to be performing well. The Department of Health through the Epidemiology Bureau takes part in achieving this goal by closely monitoring the standard surveillance indicators to ensure that the Measles elimination goal will be attained and sustained.

A total of 1,598 suspect measles-rubella cases were reported nationwide from January 1 to October 1, 2016. Of these, 1,171 were tested. Among the suspect cases, 58 (3%) were classified as confirmed measles (laboratory or epi-linked confirmed measles). One hundred seven cases (7%) were classified as laboratory confirmed rubella (Figure 1). Currently, there were no reported deaths among the confirmed measles-rubella cases.

FIGURE 1. DISTRIBUTION OF REPORTED MEASLES-RUBELLA CASES, PHILIPPINES, JANUARY 1 - OCTOBER 1, 2016 (N=1,598)




TABLE 1. MEASLES AND RUBELLA CASES BY REGION PHILIPPINES, JANUARY 1 – OCTOBER 1, 2016 (N=1,598)

REGION	REPORTED	CONFIRMED MEASLES		MEASLES COMPATIBLE	LABORATORY CONFIRMED RUBELLA	DISCARDED AS NON-MEASLES/RUBELLA	PENDING CLASSIFICATION
		LABORATORY CONFIRMED	EPI-LINKED CONFIRMED				
1	139	5	0	71	3	57	3
2	53	2	0	8	0	37	6
3	107	2	0	14	6	82	3
4A	167	6	1	33	6	111	10
4B	45	0	0	21	1	22	1
5	26	2	0	1	2	20	1
6	267	3	0	6	52	203	3
7	81	10	0	2	5	63	1
8	57	2	0	44	2	9	0
9	60	11	1	23	0	25	0
10	194	2	0	71	3	88	30
11	83	0	0	6	5	68	4
12	58	1	0	5	2	47	3
ARMM	8	1	0	3	0	1	3
CAR	70	3	0	7	7	52	1
CRG	32	2	0	12	0	18	0
NCR	151	4	0	24	13	103	7
PHL	1,598	56	2	351	107	1,006	76



TABLE 2. GEOGRAPHICAL DISTRIBUTION OF CONFIRMED CASES, PHILIPPINES, JANUARY 1 –OCTOBER 1, 2016

Measles	Region	Rubella
5	I	3
2	II	0
2	III	6
7	IVA	6
0	IVB	1
2	V	2
3	VI	52
10	VII	5
2	VIII	2
12	IX	0
2	X	3
0	XI	5
1	XII	2
1	ARMM	0
3	CAR	7
2	CARAGA	0
4	NCR	13
58	PHL	107

Map Legend:
1 DOT = 1 Case
● Measles
● Rubella





Table 2 and the maps above show the distribution of cases among regions. Twenty one percent (21%) of the confirmed measles cases came from Region IX (12 cases) while 49% of the confirmed rubella cases were from Region VI (52 cases).

**TABLE 3. NEWLY CONFIRMED CASES
MW 38 – MW 39 (n=5)**

Region	Measles	Rubella
I	1	0
IVA	1	0
IX	1	0
CAR	0	1
NCR	0	1
PHL	3	2

**TABLE 4. IDENTIFIED CLUSTERS OF CONFIRMED MEASLES AND RUBELLA
AMONG REGIONS JANUARY 1 – OCTOBER 1, 2016**

MW	Disease	Region	Province	Muncity	Barangay	Place of Transmission	No. of Cases
11	Measles	NCR	Metro Manila	Pasay City	Brgy. 46	Unknown	2
12	Measles	IX	Zamboanga del Norte	Kalawit	Palalian	Community	6
2-5	Rubella	VI	Antique	Tobias Fornier	-	School	13
9-13	Rubella	NCR	Metro Manila	Parañaque City	Tambo	Unknown	7
33	Rubella	NCR	Metro Manila	Parañaque City	Tambo	Unknown	2

No new clusters detected since MW33. Above is the list of clusters identified since January 2016.




Profile of Cases


Measles

Fifty seven percent (57%) of the confirmed measles cases were female. Majority of the confirmed cases belonged to children aged 1 to 5 years old (30%) as shown in Figure 4. Among the confirmed measles cases, 25 (46%) were not vaccinated, 24 (44%) were vaccinated and 5 (9%) have an unknown vaccination status (Figure 5).

**FIGURE 4. CONFIRMED MEASLES CASES BY AGE GROUP AND SEX
PHILIPPINES, JANUARY 1- SEPTEMBER 17, 2016 (n=54)**



**FIGURE 5. IMMUNIZATION STATUS OF CONFIRMED MEASLES CASES BY AGE GROUP
PHILIPPINES, JANUARY 1 - SEPTEMBER 17, 2016 (n=54)**






Morbidity Week 39: January 1 – October 1, 2016

Epidemiology Bureau
Public Health Surveillance Division


Among the confirmed measles cases, 24 were vaccinated with MCV. Of these, 33% received 1 dose of MCV, 21% were administered with 2 or more doses while 46% were also vaccinated but with unknown number of doses (Figure 6).

**FIGURE 6. MCV DOSE RECEIVED AMONG CONFIRMED VACCINATED MEASLES CASES
BY AGE GROUP
PHILIPPINES, JANUARY 1 - SEPTEMBER 17, 2016 (n=24)**



Out of the 54 confirmed measles cases, 25 were not vaccinated with MCV. Reasons for non-vaccination varied among the cases as shown in Figure 7.

**FIGURE 7. REASONS FOR NON-VACCINATION OF CONFIRMED MEASLES CASES
PHILIPPINES, JANUARY 1 - SEPTEMBER 17, 2016 (n=25)**






Rubella

Fifty three percent (53%) of the confirmed rubella cases were male. Majority of the confirmed cases belonged to the young adult age group ranging from 16 to 20 years old (34%) as shown in Figure 8. Among the female suspect measles-rubella cases, there were 14 pregnant cases, 2 tested positive for measles but negative for rubella and all the rest tested negative for both measles and rubella IgM. (*Pregnancy data source: CIF encoded by RITM*)

**FIGURE 8. CONFIRMED RUBELLA CASES BY AGE GROUP AND SEX
PHILIPPINES, JANUARY 1- SEPTEMBER 17, 2016 (n=101)**



Measles Surveillance Performance Indicators

**TABLE 6. MEASLES SURVEILLANCE PERFORMANCE INDICATORS* BY REGION
PHILIPPINES, 2015 vs. 2016****

REGION	POPULATION 2016	ANNUALIZED MEASLES INCIDENCE RATE		BLOOD ADEQUACY RATE		SUSPECT MEASLES CASES ADEQUATELY INVESTIGATED		ANNUALIZED SUSPECT MEASLES REPORTING RATE		ANNUALIZED NON-MEASLES/ NON-RUBELLA RATE		MEASLES COMPATIBLE %	
		Target: <1/1,000,000 Pop.		Target: ≥80%		Target: ≥80%		Target: ≥2/100,000 Pop.		Target: ≥2/100,000 Pop.		Target: <10%	
		2015	2016	2015	2016	2015	2016	2015	2016	2015	2016	2015	2016
1	5,113,827	1.58	1.04	76	50	66	47	3.27	2.95	1.96	1.30	30	49
2	3,510,762	4.04	0.76	73	83	70	81	3.61	1.78	1.33	1.33	49	15
3	11,534,111	0.71	0.23	85	84	80	76	1.40	1.17	1.00	0.90	18	14
4A	15,172,632	1.29	0.53	64	74	57	63	2.41	1.34	1.17	0.89	40	20
4B	3,057,039	1.00	0.60	47	49	38	47	2.83	1.88	0.93	0.96	55	44
5	5,920,478	0.17	0.45	96	100	92	100	0.41	0.52	0.33	0.41	13	4
6	7,703,570	8.29	0.35	97	95	89	90	5.10	4.55	3.12	3.44	7	3
7	7,565,674	6.86	1.76	98	97	86	92	2.14	1.36	1.14	1.04	5	3
8	4,430,334	4.80	0.60	20	23	16	21	4.59	1.60	0.11	0.24	87	77
9	3,814,158	28.04	3.85	70	52	62	49	10.81	1.99	4.03	0.84	35	37
10	4,865,413	13.42	0.55	43	60	42	55	6.94	5.12	0.78	2.41	69	37
11	5,033,163	25.12	0.60	92	94	87	90	6.04	2.13	2.15	1.70	20	8
12	4,768,455	17.40	0.28	60	88	57	82	8.12	1.43	2.94	1.20	42	10
ARMM	3,566,757	4.55	0.37	47	71	46	71	2.33	0.26	0.34	0.04	66	29
CAR	1,792,078	21.56	2.23	90	81	87	81	9.02	6.76	5.45	3.57	14	11
CRG	2,657,380	24.05	1.00	74	57	64	57	6.07	1.51	1.79	0.80	30	40
NCR	13,205,216	1.39	0.40	60	80	50	75	1.82	1.44	0.87	1.02	39	15
PHL	103,711,049	6.85	0.69	70	75	64	71	3.64	1.92	1.48	1.23	36	22
LEGEND:													
		<1	≥1	≥80%	<80%	≥80%	<80%	≥2/100,000 Pop.	<2/100,000 Pop.	≥2/100,000 Pop.	<2/100,000 Pop.	<10%	≤50%
													>50%



Table 6 presents the current surveillance performance of regions based on the indicators for measles surveillance. Countrywide incidence rate of 0.69 per 1,000,000 population has been achieved, reaching the target of <1 per 1,000,000 population. Regions I, VII, IX, CAR and CARAGA did not meet the target which implies increased occurrence of measles in these regions.

These surveillance indicators gauges the capacity of the country in achieving the measles elimination goal. Analyzing the overall performance of all the surveillance indicators, the country needs a joint effort among regions in order to cope up with these targets. Intensification of active surveillance should be initiated across the country in order to reach the targets towards measles elimination.

Annex A. Definition of Terms

Laboratory confirmed measles case	☞ A suspect measles case with a positive laboratory test result for measles-specific IgM antibodies or other approved laboratory test method
Laboratory confirmed rubella case	☞ A suspect measles case with a positive laboratory test result for rubella-specific IgM antibodies or other approved laboratory test method
Measles compatible case	☞ A case that meets the suspect case definition for measles but for which no adequate blood specimen was taken and which has not been linked epidemiologically to another case positive for measles IgM or another laboratory-confirmed communicable disease
Confirmed Measles cases	☞ Laboratory confirmed + Epidemiologically-linked measles cases
Epidemiologically-linked measles (or rubella) case	☞ A suspect measles case that has not been confirmed by laboratory but that is geographically AND temporally related (with dates of rash onset occurring between 7 and 21 days apart) to a laboratory-confirmed case or (in the event of an outbreak) to another epidemiologically confirmed measles case.
Discarded as non-measles/non-rubella	☞ A case that meets the clinical case definition for measles and discarded as non-measles/rubella case.
Pending Classification	☞ Cases with blood specimen collected and pending laboratory results.
Alert threshold	☞ Refers to the level of occurrence of disease that serves as an early warning for epidemics. An increase in the number of cases above the threshold level should trigger an investigation, epidemic preparedness and implement appropriate prevention and control measures.
Epidemic threshold	☞ Refers to the level of occurrence of disease above which an urgent response is required. The threshold is specific to each disease and depends on the infectiousness, other determinants of transmission and local endemicity levels.
Cluster of cases	☞ 2 or more cases with temporal (occurring in a span of 4 weeks) and geographical association (within the same barangay)

Annex B. Measles Surveillance Indicators Targets

Measles incidence rate*, target: <1/ 1,000,000 of the total population. It measures the progress of a country towards measles elimination. High incidence rate indicates persistence of measles transmission in some areas.

Suspect Measles Reporting Rate (or Measles Rate)*, target: ≥2 per 100,000 of the total population. It measures the ability to detect suspect measles cases. Reporting an adequate number of suspected cases provides confidence that the system is sensitive to detect measles cases.



Morbidity Week 39: January 1 – October 1, 2016

Epidemiology Bureau
Public Health Surveillance Division

Non-Measles Reporting Rate*, target: ≥ 2 per 100,000 of the total population. If non-measles reporting rate is equal or proportion to the number of suspected measles cases in all regions, it gives us higher chance in attaining our goal of measles elimination.

Adequacy of blood specimen (blood adequacy rate), target: $\geq 80\%$ adequate specimen collection rate. This will facilitate the specificity (ability to determine measles virus as the cause of illness) of reported measles cases. With adequate specimen collection there will be an access to identify the circulating measles virus in the community.

Timeliness and adequacy of investigation, provides venue to prevent further transmission of measles cases in the community, furthermore, provides immediate response to prevent potential outbreaks. Its target rate is $\geq 80\%$ of cases were investigated within 48 hours of notification.

*Annualized rate, measures the incidence or reporting in a period of 1 year. This is computed by the number of specific measles cases over the target measles cases divided by 12 months then multiplied by the number of months to be analyzed.

Editorial Board


IRMA L. ASUNCION, MD, MHA, CESO IV
Director IV, Epidemiology Bureau


GENESIS MAY J. SAMONTE, MD, PHSAE
OIC-Division Chief, PHSD


JUNE CANTATA B. CORPUZ, RN
Nurse III, PIDSR National Coordinator


JEZZA JONAH D. CRUCENA, RN
Nurse III, VPDS National Coordinator


ALLAN P. IGNACIO
Statistician II, PHSD

